


AMENDMENTS TO THE CLAIMS

Please amend claims 1, 7, 17, 20, 21, 23-29, 31, and 32 as indicated below.
Deletions appear in ~~strikethrough font~~, and additions are underlined.

Complete listing of claims

1. **(Currently amended)** A substituted heterocyclo-norbornylamino ~~derivative~~
compound having an exo-configuration nitrogen and endo-fused five-membered
or six-membered ring of the formula I or having an exo-configuration nitrogen and
exo-fused five-membered or six-membered ring of the formula Ia,



wherein:

A is (C₁-C₄)- alkylene;

T is (C₁-C₄)- alkyl or H;

B is a saturated or unsaturated carbon five-membered or six-membered ring, which is unsubstituted or is substituted by 1-3 substituents chosen from oxo, hydroxyl, (C₁-C₄)- alkoxy, and (C₁-C₄)- alkyl;

Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to four identical or different heteroatoms chosen from O, S, N, and Se;

R is OH, F, Cl, Br, I, CN, NO₂, phenyl, CO₂R₁, (C₁-C₄)- alkyl, (C₁-C₄)-alkoxy, amino, (C₁-C₄)- alkylamino, di(C₁-C₄)- alkylamino, or amino-(C₁-C₄)- alkyl,

wherein the alkyl radicals are unsubstituted or are completely or partly substituted by fluorine;

R₁ is H or (C₁-C₄)-alkyl, which is unsubstituted or completely or partly substituted by fluorine;

n is 0, 1, 2, 3 or 4,

wherein, if n = 2, 3 or 4, the substituents R are chosen independently of one another;

or a pharmaceutically tolerable salt or trifluoroacetate thereof.

2. **(Original)** A compound as claimed in claim 1, wherein:

A is (C₁-C₂)- alkylene;

T is H or methyl;

B is a saturated or unsaturated carbon five-membered or six-membered ring;

Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to three identical or different heteroatoms chosen from O, S, and N;

R is F, Cl, Br, iodine, amino, hydroxymethyl, OH, phenyl, CO₂R₁, (C₁-C₄)-alkyl, or (C₁-C₄)- alkoxy,

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

wherein the alkyl radicals are unsubstituted or completely or partly substituted by fluorine;

R₁ is H or (C₁-C₄)-alkyl, where the alkyl radical is unsubstituted or completely or partly substituted by fluorine;

n is 0, 1, 2 or 3,

where, if n = 2 or 3, the corresponding substituents R are chosen independently of one another.

3. **(Original)** A compound as claimed in claim 1, wherein:

A is (C₁-C₂)-alkylene;

T is hydrogen;

B is a saturated or unsaturated carbon five-membered or six-membered ring;

Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to two identical or different heteroatoms chosen from O, S, and N ;

R is F, Cl, Br, (C₁-C₄)-alkoxy, or (C₁-C₄)-alkyl,
where the alkyl radicals are unsubstituted or completely or partly substituted by fluorine;

n is 0, 1 or 2, wherein, if n = 2, the corresponding substituents R are chosen independently of one another.

4. **(Original)** A compound as claimed in claim 1, chosen from:

exo/exo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,

(rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
(+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
(-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
(rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
(+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
(-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)pyrazin-2-ylmethylamine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-2-ylmethylamine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-3-ylmethylamine,
exo/endo-(3a,4,5,6,7,7a-hexahydro-3H-4,7-methanoinden-5-yl)pyridin-3-yl-
methylamine,
exo/endo-(3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-yl)pyridin-3-yl-
methylamine,
exo/endo-furan-3-ylmethyl-(octahydro-4,7-methanoinden-5-yl)amine,
exo/endo-furan-2-ylmethyl-(octahydro-4,7-methanoinden-5-yl)amine,
exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)pyridin-3-ylmethylamine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)-(1H-pyrrol-2-ylmethyl)amine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)-pyrimidin-5-ylmethylamine, and
their pharmaceutically tolerable salts or trifluoroacetates.

5. **(Original)** A compound as claimed in claim 1, chosen from:

exo/exo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
(rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
(+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP



1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com

(-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
(rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
(+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-2-ylmethylamine,
exo/endo-(3a,4,5,6,7,7a-hexahydro-3H-4,7-methanoinden-5-yl)pyridin-3-yl-
methylamine,
exo/endo-(3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-yl)pyridin-3-yl-
methylamine,

exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)pyridin-3-ylmethylamine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)-(1H-pyrrol-2-ylmethyl)amine,
exo/endo-(octahydro-4,7-methanoinden-5-yl)-pyrimidin-5-ylmethylamine, and
their pharmaceutically tolerable salts or trifluoroacetates.

6. **(Original)** A process for the preparation of a compound as claimed in claim 1,
comprising


reacting a compound of the formula (II) or (II a)



with a compound of the formula (III)

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER LLP

1300 I Street, NW
Washington, DC 20005
202.408.4000
Fax 202.408.4400
www.finnegan.com




in the presence of at least one reductant or at least one Lewis acid to give at least one compound of the formula (I) or (I a),



wherein, independently of one another, A' corresponds to a bond or (C₁-C₃)-alkyl, A'' corresponds to H or (C₁-C₃)-alkyl, and A' and A'' together with the carbon atom of the carbonyl group represent as many carbon atoms as A represents in formula (I) or (I a); and

optionally converting the compound of formula (I) or (I a) into a pharmaceutically tolerable salt or trifluoroacetate.


7. **(Currently amended)** A process for the preparation of a compound as claimed in claim 1, comprising
isolating an intermediate of the formula (IV) or (IV a),



formed from reacting a compound of the formula (II) or (II a)



with a compound of the formula (III),



and then converting the intermediate of the formula (IV) or (IV a) into a compound of the formula (I) or (I a) by using at least one reductant,

wherein, independently of one another, A' corresponds to a bond or (C_1-C_3) -alkyl, A'' corresponds to H or (C_1-C_3) -alkyl, and A' and A'' together with the carbon atom of the carbonyl group represent as many carbon atoms as A represents in formula (I) or (I a);



wherein T# is a free electron pair or (C_1-C_4) -alkyl, and

wherein a counterion is assigned to the iminium ion formed when T# is (C_1-C_4) -alkyl; and


optionally converting the compound of formula (I) or (I a) into a pharmaceutically tolerable salt or trifluoroacetate.

8. **(Original)** A process as claimed in claim 7, wherein the counterion assigned to the ammonium nitrogen formed when T# is (C_1-C_4) -alkyl is chosen from chloride and tosylate.

9. **(Original)** A process for the preparation of a compound as claimed in claim 1, comprising reacting a compound of the formula (II) or (II a)



with an alkylating agent of the formula (V) to produce a compound of the formula (I) or (I a),




wherein U is a nucleophilically substitutable group and wherein, independently of one another, A' corresponds to a bond or (C_1-C_3) -alkyl, A'' corresponds to H or (C_1-C_3) -alkyl, and A' and A'' together with the carbon atom which U is bonded represent as many carbon atoms as A represents in formula (I) or (I a); and

optionally converting the compound of formula (I) or (I a) into a pharmaceutically tolerable salt or trifluoroacetate.

10. **(Original)** A process as claimed in claim 9, wherein U is chosen from halogen atoms, alkylsulfonates, and arylsulfonates.

11. **(Original)** A process as claimed in claim 10, wherein U is chosen from Cl, Br, I, mesylate, and tosylate.
12. **(Original)** A process for the preparation of a compound as claimed in claim 1, comprising reducing carboxamides of the formula (VI) or (VI a) to the corresponding amines of the formula (I) or (I a),



wherein A* corresponds to a bond or (C₁-C₃)-alkyl, and

optionally converting the compound of formula (I) or (I a) into a pharmaceutically tolerable salt or trifluoroacetate.

13. **(Original)** A process for the preparation of a compound as claimed in claim 1, comprising alkylating a compound of the formula (I) or (I a), in which T corresponds to hydrogen, using alkylating agents of the formula (VII),




wherein T* is (C₁-C₄)-alkyl and U is a nucleophilically substitutable group,


such that tertiary amines result from this reaction; and

optionally converting the compound of formula (I) or (I a) into a pharmaceutically tolerable salt or trifluoroacetate.

14. **(Original)** A process as claimed in claim 13, wherein U is chosen from halogen atoms, alkylsulfonates, and arylsulfonates.
15. **(Original)** A process as claimed in claim 14, wherein U is chosen from Cl, Br, I, mesylate, and tosylate.
16. **(Original)** A process for the preparation of a compound as claimed in claim 1, comprising reacting a dicyclopentadienylplatinum complex of the formula (VIII)



with amines of the formula (IX),



and subsequently reducing the intermediate formed to a compound of the formula (I) or (I a)

wherein, independently of one another, A' corresponds to a bond or (C₁-C₃)-alkyl, A'' corresponds to H or (C₁-C₃)-alkyl, and A' and A'', together with the carbon atom to which the nitrogen atom is bonded, represent as many carbon atoms as A represents in formula (I), and

optionally converting the compound of formula (I) into a pharmaceutically tolerable salt or trifluoroacetate.

17. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of a disorder of the respiratory drive, comprising administering to a patient in need thereof a compound as claimed in claim 1.
18. **(Original)** A method according to claim 17, wherein the disorder of the respiratory drive is a sleep-related respiratory disorder.
19. **(Original)** A method according to claim 18, wherein the sleep-related respiratory disorder is sleep apnea.
20. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of snoring, comprising administering to a patient in need thereof a compound as claimed in claim 1.
21. **(Currently amended)** A method for the treatment ~~or for the prophylaxis~~ of acute or chronic kidney disorders, comprising administering to a patient in need thereof a compound as claimed in claim 1.

22. **(Original)** A method according to claim 21, wherein the acute or chronic kidney disorder is chosen from acute kidney failure and chronic kidney failure.
23. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of disorders of the intestinal function, comprising administering to a patient in need thereof a compound as claimed in claim 1.
24. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of disorders of the bile function, comprising administering to a patient in need thereof a compound as claimed in claim 1.
25. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of an ischemic condition of the peripheral or central nervous system, or an ischemic condition due to a stroke, comprising administering to a patient in need thereof a compound as claimed in claim 1.
26. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of an ischemic condition of a peripheral organ or a limb, comprising administering to a patient in need thereof a compound as claimed in claim 1.
27. **(Currently amended)** A method for the treatment of a an ischemia-induced endogenous state of shock, comprising administering to a patient in need thereof a compound as claimed in claim 1.
28. **(Currently amended)** A method for the protection of an organ during use in a surgical operation or an organ transplantation, comprising administering to a patient in need thereof a compound as claimed in claim 1.
29. **(Currently amended)** A method for the conservation or storage of an organ to be transplanted ~~a transplant for surgical measures, comprising administering to a~~

~~patient in need thereof a~~ storing said organ in a solution comprising a compound
as claimed in claim 1.

30. **(Original)** A method for the treatment of a disease in which cell proliferation is a primary or secondary cause, comprising administering to a patient in need thereof a compound as claimed in claim 1.

31. **(Currently amended)** A method for the treatment ~~or prophylaxis~~ of a disorder of lipid metabolism, comprising administering to a patient in need thereof a compound as claimed in claim 1.

32. **(Currently amended)** A method for the treatment ~~or prophylaxis of attack of an~~ infestation by an ectoparasite, comprising administering to a patient in need thereof a compound as claimed in claim 1.

33. **(Original)** A composition, comprising at least one compound as claimed in claim 1 and a carrier.

34. **(Original)** A pharmaceutical composition, comprising at least one compound as claimed in claim 1 and a pharmaceutically acceptable carrier.